

General Canning Information

Identifying and Handling Spoiled Canned Food

Do not taste food from a jar with an unsealed lid or food that shows signs of spoilage. You can more easily detect some types of spoilage in jars stored without screw bands. Growth of spoilage bacteria and yeast produces gas which pressurizes the food, swells lids, and breaks jar seals. As each stored jar is selected for use, examine its lid for tightness and vacuum. Lids with concave centers have good seals.

Next, while holding the jar upright at eye level, rotate the jar and examine its outside surface for streaks of dried food originating at the top of the jar. Look at the contents for rising air bubbles and unnatural color.

While opening the jar, smell for unnatural odors and look for spurting liquid and cotton-like mold growth (white, blue, black, or green) on the top food surface and underside of lid.

Spoiled low-acid foods, including tomatoes, may exhibit different kinds of spoilage evidence or very little evidence. Therefore, all suspect containers of spoiled low-acid foods, including tomatoes, should be treated as having produced botulinum toxin and handled carefully in one of two ways:

- If the swollen metal cans or suspect glass jars are still sealed, place them in a heavy garbage bag. Close and place the bag in a regular trash container or dispose in a nearby landfill.
- If the suspect cans or glass jars are unsealed, open, or leaking, they should be detoxified before disposal.

Detoxification process: Wear disposable rubber or heavy plastic gloves.

Carefully place the suspect containers and lids on their sides in an 8-quart volume or larger stock pot, pan, or boiling-water canner. Wash your hands with gloves thoroughly. Carefully add water to the pot and avoid splashing the water. The water should completely cover the containers with a minimum of a 1-inch level above the containers. Place a lid on the pot and heat the water to boiling. Boil 30 minutes to ensure detoxifying the food and all container components. Cool and discard the containers, their lids, and food in the trash or dispose in a nearby landfill.

Cleaning up the area: Contact with botulinum toxin can be fatal whether it is ingested or enters through the skin. Take care to avoid contact with suspect foods or liquids. Wear rubber or heavy plastic gloves when handling suspect foods or

cleaning up contaminated work surfaces and equipment. A fresh solution of 1 part unscented liquid household chlorine bleach (5 to 6% sodium hypochlorite) to 5 parts clean water should be used to treat work surfaces, equipment, or other items, including can openers and clothing, that may have come in contact with suspect foods or liquids. Spray or wet contaminated surfaces with the bleach solution and let stand for 30 minutes. Wearing gloves, wipe up treated spills with paper towels being careful to minimize the spread of contamination. Dispose of these paper towels by placing them in a plastic bag before putting them in the trash. Next, apply the bleach solution to all surfaces and equipment again, and let stand for 30 minutes and rinse. As a last step, thoroughly wash all detoxified counters, containers, equipment, clothing, etc. Discard gloves when cleaning process is complete. (Note: Bleach is an irritant itself and should not be inhaled or allowed to come in contact with the skin.)

This document was adapted from the *"Complete Guide to Home Canning,"* Agriculture Information Bulletin No. 539, USDA, revised 2009.

Reviewed November 2009.